

# Rely+On<sup>™</sup> PeraSafe<sup>™</sup>

High Level Disinfectant for use with Medical Devices



- · Outstanding biocidal action
- Optimised corrosion inhibitor system
- Suitable for the disinfection of endoscopes, ultra-sound probes and other reusable, heat sensitive medical devices
- · Good environmental and operator safety profiles



### Rely+On<sup>™</sup> PeraSafe<sup>™</sup>

Specifically developed to offer rapid broad spectrum decontamination against viruses, mycobacteria, bacteria, yeasts and bacterial spores. Rely+On<sup>TM</sup> PeraSafe<sup>TM</sup>, with its optimised corrosion inhibitor system and near neutral pH, is a suitable solution for endoscope, ultra-sound probe and other reusable, heat sensitive medical device disinfection.

### Biocidal Efficacy

Rely+On<sup>TM</sup> PeraSafe<sup>TM</sup> powder generates peracetic acid *in-situ* when dissolved in water. The neutral pH solution combines the biocidal efficacy of peracetic acid with excellent materials compatibility to allow repeated, rapid decontamination of pathogenic micro-organisms from heat labile, reusable medical devices and instruments.

#### **Proven Broad Spectrum Efficacy**

Independently proven highly effective against viruses, mycobacteria, bacteria and bacterial spores, Rely+On<sup>TM</sup> PeraSafe<sup>TM</sup> provides rapid broad spectrum decontamination against disease-causing pathogens of concern to human health, such as, HIV, Hepatitis B & C, Mycobacteria (Tuberculosis),Pseudomonas aeruginosa, Clostridium difficile, MRSA and Candida albicans.

Rely+On™ PeraSafe™ has been independently evaluated in the United Kingdom by the Hospital Infection Research Laboratory (HIRL).

The HIRL evaluation findings demonstrate that, in 10 minutes or less, Rely+On™ PeraSafe™ reduces colony forming units of >5 log<sub>10</sub> for Bacillus subtilis var niger spores (surface test), and >5 log<sub>10</sub> for Mycobacterium terrae. These results clearly demonstrate Rely+On™ PeraSafe™'s rapid biocidal action and suitability for use in medical device disinfection.

#### **Medical Device Compatibility**

Rely+On<sup>TM</sup> PeraSafe<sup>TM</sup> is compatible with endoscopes, ultra-sound probes, surgical instruments and other common reusable, heat sensitive medical devices. An optimised corrosion inhibitor system, allied to a neutral pH solution, optimises efficacy and compatibility.



#### Certification

Rely+On<sup>™</sup> PeraSafe<sup>™</sup> complies with the Medical Devices Directive as assessed by BSI Product Services (Notified Body No. 0086) (CE Certificate No. 01455).



#### **Operator Safety**

Rely+On<sup>™</sup> PeraSafe<sup>™</sup> is not classified as a sensitiser in both the powdered form, and the in-use dilutions.

#### **Environmental Profile**

Rely+On™ PeraSafe™ has been formulated to include ingredients that have been carefully selected for their ability to degrade naturally within the environment.

#### **Lifespan of Disinfectant Solution**

Maximum 24 hours from activation and/or 20 immersions, whichever is the least. For fungicidal activity claims, solutions are to be used within 8 hours of preparation. Unused or inactive solutions may be disposed of via the normal waste water discharge channels\*.

## Rely+On™ PeraSafe™ is available in the following presentations:

Pack Size	Volume of disinfectant solution
16.2 gram (sachet)	1 litre
81 gram (bottle)	5 litres
162 gram (bottle)	10 litres
810 gram (bottle)	50 litres

\* Always check and observe the local site requirements concerning the disposal of solutions entering waste water treatment facilities.



#### **Easy to Prepare**



**Step 1**Fill a container with the appropriate volume of lukewarm tap water (30-35°C).



Step 2
Prepare the
solution by adding
81 grams of
Rely+On™ PeraSafe™
powder to every
5 litres of water.



Step 3
Stir until the powder dissolves to form a blue solution. Then leave for 15 minutes to allow the disinfectant to fully activate.

The tables below summarise independent efficacy data of Rely+On™ PeraSafe™ against the important disease-causing organisms affecting human health.

#### Virucidal Efficacy

Organism	Strain	Test Protocol	Contact Time	Log Reduction
Adenovirus	Type 5	EN 14476	5 minutes	>4 log
Hepatitis B	DHBV strain	EN 14476	10 minutes	>4 log
Hepatitis C	BVD surrogate strain	EN 14476	10 minutes	>4 log
HIV	HTLV-IIIB	USA EPA 40 part 158	5 minutes	>4 log
Poliovirus	Type 1	EN 14476	5 minutes	>4 log
Murine norovirus	s99	EN 14476	10 minutes	>4 log

#### Mycobactericidal Efficacy

Organism	Strain	Test Protocol	Contact Time	Log Reduction
Mycobacterium terrae	ATCC 15769	EN 14563	10 minutes	>4 log
Mycobacterium avium	ATCC 15755	EN 14563	10 minutes	>4 log
Mycobacterium terrae	NCTC 10856	EN 14348	10 minutes	>5 log

### Bactericidal Efficacy

Organism	Strain	Test Protocol	Contact Time	Log Reduction
Campylobacter jejuni	Wild strain	EN 1276	5 minutes	>5 log
Escherichia coli	NCTC 12900	EN 1276	5 minutes	>5 log
Enterococcus hirae	NCTC 12367	EN 13727	10 minutes	>5 log
Enterococcus hirae	ATCC 10541	EN 14561	10 minutes	>5 log
Klebsiella aerogenes	NCTC 9528	EN 1276	5 minutes	>5 log
Staphylococcus aureus	MRSA NCTC 12493	EN 1276	5 minutes	>5 log
Staphylococcus aureus	EMRSA type 15	EN 1276	5 minutes	>5 log
Staphylococcus aureus	NCTC 10788	EN 13727	10 minutes	>5 log
Staphylococcus aureus	ATCC 6538	EN 14561	10 minutes	>5 log
Salmonella typhimurium	Phage type 104	EN 1276	5 minutes	>5 log
Pseudomonas aeruginosa	NCTC 6749	Endoscope in-use trial	4 minutes	>7 log
Pseudomonas aeruginosa	NCTC 6749	EN 13727	10 minutes	>5 log
Pseudomonas aeruginosa	ATCC 15442	EN 14561	10 minutes	>5 log

#### **Bactericidal Spore Efficacy**

Organism	Strain	Test Protocol	Contact Time	Log Reduction
Bacillus subtilis var niger (spores)	NCTC 10073	EN 13704 (modified)	5 minutes	>4 log
Clostridium difficile	Stool isolate	Surface	10 minutes	>6 log

#### Fungicidal / Yeasticidal Efficacy

Organism/Disease	Strain	Test Protocol	Contact Time	Log Reduction
Aspergillus niger (spores)	ATCC 16404	EN 13624	60 minutes*	>4 log
Aspergillus niger (spores)	ATCC 16404	EN 14562	30 minutes	>4 log
Candida albicans	ATCC 10231	EN 13624	10 minutes	>4 log
Candida albicans	ATCC 10231	EN 13624	60 minutes*	>4 log
Candida albicans	ATCC 10231	EN 14562	30 minutes	>4 log

Full copies of the independent test data summarised above are available upon request from Lanxess. \*8 hour solution data. \*\*Fungicidal claims for the product are only valid for solutions up to 8 hours old, from the time of first preparation.



Antec International Limited
Windham Road, Chilton Industrial Estate
Sudbury, Suffolk CO10 2XD, United Kingdom

Tel: +44(0)1787 377305 relyondisinfection@lanxess.com www.relyondisinfection.com